

Determining the expansion history of the universe with the three dimensional cosmic shear

Ram Brustein and Daniel Levy

Department of Physics, Ben-Gurion University, Beer-Sheva 84105, Israel

e-mail: ramyb@bgu.ac.il, levyda@bgu.ac.il

We define a new observable that depends on finite redshift differences of the spin-weighted angular moments of the two point function of the three dimensional cosmic shear and on luminosity distance. It is shown that precise measurements of our observable will be able to tightly constrain the expansion history of the universe.

The observational evidence that the expansion of the universe is being accelerated by dark energy (DE) is strengthening continuously [1, 2, 3, 4, 5]. On the other hand, the theoretical efforts to understand the nature of DE and the underlying reason for the accelerated expansion were less successful so far. It has become clear that in order to make further theoretical progress better methods of determining the expansion history of the universe are needed. The kinematic distance probes in the homogeneous and isotropic universe, such as luminosity distance (d_L), angular distance etc., are limited because they rely on light emitted from distant sources, and hence measure an integral over the expansion history [6]. It is possible to circumvent some of their limitations by adding prior assumptions on the evolution, or by combining several kinds of kinematic observations [7, 8, 9], or by focusing on better determined quantities [10, 11]. A recent review of the subject can be found in [12].

In addition to the kinematic probes one can use dynamical (fluctuations) probes to determine the expansion history of the universe. Cosmological perturbations provide, through their dependence on the homogeneous and isotropic background, an independent source of information [13, 14]. One needs observables that can determine in a reliable way the time dependence of the perturbations' amplitude and not only their spectrum at a fixed redshift. Such observables can have a simple functional dependence on the expansion history and hence can provide much more information than a measurement of a single spectrum. However, to provide significant constraints above and beyond those that kinematical probes put on the expansion history such observables have to be measured and evaluated precisely.

The three dimensional cosmic shear (3DCS) is a redshift dependent tensor that measures the shape distortion of distant galaxies as the light emitted from them propagates through the perturbed universe [16]. The two dimensional cosmic shear was recently measured in weak lensing (WL) experiments [17, 18, 19] while planned programs are expected to have some three dimensional capabilities [20]. Several recent reviews of the theory and observations of WL provide an excellent description of the field [27, 28]. Our results suggest that precise measurements of the 3DCS will significantly enhance our ability to constrain the expansion history of the universe.

If the cosmological perturbations are small, the cosmic shear (CS) depends linearly on them. In this linear regime both observations and theoretical calculations can be done accurately. However, to take full advantage of the linear regime one needs to filter the influence of the late time nonlinear regime [29]. This requires the capability of accurately measuring finite differences of the 3DCS at different redshifts.

The calculation of the 3DCS relies on solving the geodesic deviation equation in the presence of the cosmological perturbations to calculate the rate of divergence of light propagating on nearby geodesics. Obviously, this requires knowing the rate of divergence of light propagating on nearby geodesics in the absence of the perturbations, which gives the angular distance in the unperturbed universe. Hence accurate knowledge of the angular distance (or equivalently, d_L) is necessary for accurate knowledge of the 3DCS.

In this paper we define a new observable $J(z)$ derived from the 3DCS, and discuss how precise measurements of J can be used to tightly constrain the expansion history of the universe. The observable $J(z)$ depends on finite redshift differences of the spin-weighted angular moments of the two point function of the 3DCS. The equations connecting the metric perturbations and the CS that allow us to make the quantitative estimates in this paper were developed in [30].

We consider the perturbed spatially flat Friedman-Robertson-Walker universe. We assume that the cosmic fluid is a perfect fluid and that the perturbations are adiabatic. In this case the line element of the perturbed universe is defined in terms of single space-time dependent perturbation Φ

$$ds^2 = a^2(\eta) \left[-(1 + \Phi(\eta, \vec{x})) d\eta^2 + (1 - \Phi(\eta, \vec{x})) (dw^2 + w^2 d\Omega^2) \right]. \quad (1)$$

Here η is conformal time and w is a radial coordinate. In [30] we have followed the standard

derivations that are reviewed in [31] to obtain the equations of motion for Φ in terms of the total redshift dependent equation-of-state (EOS) parameter $w_{\text{tot}}(z) = p(z)/\rho(z)$.

We briefly review the results of [30]. When the universe is matter dominated the growing mode of Φ factorizes $\Phi_+(z, \vec{x}) = C(\vec{x})\Phi_T(z)$. The equation for Φ_T in terms of w_{tot}

$$\Phi_T(z) - \frac{2(1+z)}{5+3w_{\text{tot}}(z)}\partial_z\Phi_T(z) = \frac{1+w_{\text{tot}}(z)}{5+3w_{\text{tot}}(z)} \quad (2)$$

can be solved as follows. First, define an integration factor

$$I(z) = \int_z^{z_{\text{in}}} d\tilde{z} \frac{5+3w_{\text{tot}}(\tilde{z})}{2(1+\tilde{z})}. \quad (3)$$

Then reduce eq.(2) to a first order equation

$$e^{-I(z)}\partial_z [e^{I(z)}\Phi_T(z)] = -\frac{1+w_{\text{tot}}(z)}{2(1+z)} \quad (4)$$

whose solution is

$$\Phi_T(z) = \Phi_T(z_{\text{in}}) + e^{-I(z)} \int_z^{z_{\text{in}}} d\tilde{z} e^{I(\tilde{z})} \frac{1+w_{\text{tot}}(\tilde{z})}{2(1+\tilde{z})}. \quad (5)$$

The initial conditions are set at the initial redshift z_{in} . Asymptotically, for very large z the universe was matter dominated, hence $w_{\text{tot}} \rightarrow 0$ and Φ_T is a constant whose value can be determine from eq.(2): $\Phi_T \rightarrow 1/5$.

We assume a power-law spectrum for Φ_+ . In momentum space this implies that $q^3\langle C(\vec{q}_1), C^*(\vec{q}_2) \rangle = A(q/q_0)^{n-1}\delta(\vec{q}_1 - \vec{q}_2)$ where n is the spectral index and A is the spectral amplitude at q_0 . Both were measured most recently by the WMAP and SDSS experiments [3, 4]. In particular, the spectral index is approximately $n = .95$ corresponding to an approximately flat spectrum. The range over which the spectrum is flat (n is approximately 1) is limited because causal processes inside the horizon suppress the perturbations [32]. (For further discussion see [30]). In the rest of the paper we will assume for simplicity a flat spectrum $n = 1$. This assumption can be relaxed [33].

The CS tensor γ quantifies the distortion pattern of distant objects as light emitted from them propagates through the perturbed universe. It can be measured in WL observations. We will be interested in the angular moments of γ . Because γ is a tensor it has to be expanded in tensor spherical harmonics $\gamma = \sum_{lm} {}_2a_{lm} {}_2Y_{lm}$, and the two-point correlation function of γ has a similar expression $\langle \gamma, \gamma^* \rangle = \sum_l \sqrt{\frac{2l+1}{4\pi}} {}_2C_l {}_2Y_{l,-2}$. The relevant properties

of the s -weighted spherical harmonics ${}_sY_{l,m}$ can be found in [34]. The spin-weight operator $\bar{\partial}$ operates on the s -weight spherical harmonic and gives an $s + 1$ -weight spherical harmonic $\bar{\partial} {}_sY_{l,m} = [(l - s)(l + s + 1)]^{1/2} {}_{s+1}Y_{l,m}$.

In [30] we have shown that

$$\gamma(w, \theta, \varphi) = \int_0^w dw' \frac{(w - w')}{w w'} \bar{\partial} \bar{\partial} \Phi(w', \theta, \varphi) \quad (6)$$

and that the angular coefficients ${}_2C_l$ are consequently

$${}_2C_l(w_1, w_2) = (4\pi)^2 Al(l + 2) \int_0^{w_1} du \frac{(w_1 - u)(w_2 - u)}{u w_1 w_2} \Phi_T^2(u) \quad , \quad w_1 < w_2. \quad (7)$$

We now define the new observable $J(z)$. As we have explained, to take full advantage of the linear regime the observable has to subtract the region of small u in eq.(7) where the integral formally diverges. However, the subtraction has to be of a kind that will allow a good measurement of the defined quantity. After considering different possibilities we have come up with the simple finite difference

$$\Delta_{w_2} {}_2C_l(w_1, w_2) = \frac{1}{(4\pi)^2 Al(l + 2)} \left({}_2C_l(w_1, w_2 + \Delta w_2) - {}_2C_l(w_1, w_2) \right). \quad (8)$$

Evaluating $\Delta_{w_2} {}_2C_l(w_1, w_2)$ using eq.(7) we obtain

$$\Delta_{w_2} {}_2C_l(w_1, w_2) = \frac{1}{w_1 w_2} \frac{\Delta w_2}{w_2 + \Delta w_2} \int_0^{w_1} du (w_1 - u) \Phi_T^2(u). \quad (9)$$

This suggests the definition

$$J(w) = \int_0^w du (w - u) \Phi_T^2(u). \quad (10)$$

Since $J(w)$ is l , w_2 and Δw_2 -independent it can be estimated by statistically averaging measurements for many different l 's and many different values of w_2 and Δw_2 ,

$$J(w) = \left\langle w w_2 \frac{w_2 + \Delta w_2}{\Delta w_2} \Delta_{w_2} {}_2C_l(w, w_2) \right\rangle_{l, w_2, \Delta w_2}, \quad (11)$$

the only restriction is that $w_2 > w$.

Since $w(z) = \int_0^z \frac{dy}{H(y)} = \frac{d_L}{1+z}$ we may express J as a function of redshift

$$J(z) = \int_0^z dy \frac{1}{H(y)} \left(\frac{d_L(z)}{1+z} - \frac{d_L(y)}{1+y} \right) \Phi_T^2(y). \quad (12)$$

We would like to show that accurate measurements of J can significantly enhance our ability to constrain the expansion history of the universe. We will use eq.(11) to determine the requirements for an accurate measurement of J . First one has to measure accurately the angular moments ${}_2C_l(w_1, w_2)$ for many l 's. Since the measurements of the CS are noisy because of intrinsic ellipticity variations and observational limitations one uses statistical methods to reduce the errors. The more galaxies are measured, the better the shear measurement. The best that one can hope to do is to measure the angular moments such that the errors are dominated by statistics. Hence, the ultimate limitation is set by cosmic variance. By averaging over many angular moments from l_1 to l_2 ($l_1, l_2 \gg 1$) one can reach a relative accuracy of $\sim 1/\sqrt{l_1^2 - l_2^2}$. The largest l for which the ${}_2C_l$'s can be measured accurately and the averaging can be done reliably will determine the ultimate accuracy. Thus, to get below 1% accuracy one would need to make a cosmic variance limited measurement of about a 100 angular moments in the range $l \sim 100$.

Precise measurements of the angular moments are not sufficient. In addition $w(z)$ and $\Delta w(z)$ have to be measured precisely. This means that the redshift at which the angular moments are measured and d_L for this redshift have to be known precisely. If one wishes to obtain a 1% measurement of J then redshift measurements and d_L measurement at or below 1% have to be made. While a 1% redshift measurement is significantly below the expected accuracy of photometric redshift measurement it can, of course, be measured to much better accuracy using spectroscopic methods. A 1% measurement of d_L is a plausible goal for future supernova observations, however, accuracies of much below 1% are not likely. Hence the final limiting factor on the accuracy of measuring J is likely to be set by the accuracy of d_L measurements. To summarize, measuring J with one percent precision (or perhaps somewhat better) seems a plausible optimistic goal, though a great challenge for future observations.

We wish to estimate the sensitivity $\delta J/\delta w_{\text{tot}}$. It is very instructive to obtain first the analytical the dependence of the sensitivity on different parameters. We start by calculating

$\delta\Phi_T/\delta w_{\text{tot}}$. From eq.(2) we find after some algebra that

$$\begin{aligned} e^{I(z)} \left(\Phi_T(z) \int_z^{z_i} d\tilde{z} \frac{3\delta w_{\text{tot}}(\tilde{z})}{2(1+\tilde{z})} + \delta\Phi_T(z) \right) - e^{I(z_i)} \delta\Phi_T(z_i) = \\ = \int_z^{z_i} d\tilde{z} \frac{e^{I(\tilde{z})}}{2(1+\tilde{z})} \left[(1+w_{\text{tot}}(\tilde{z})) \int_{\tilde{z}}^{z_i} dx \frac{3\delta w_{\text{tot}}(x)}{2(1+x)} + \delta w_{\text{tot}}(\tilde{z}) \right]. \end{aligned} \quad (13)$$

Equation (13) determines $\delta\Phi_T$ in terms of δw_{tot} .

We may gain more insight by restricting to a specific case $\delta w_{\text{tot}} = \begin{cases} \epsilon & z_1 < z < z_2 \\ 0 & \text{Otherwise} \end{cases}$. In

this case we may choose the upper limit to be $z_i = z_2$. To calculate $\delta\Phi/\delta w_{\text{tot}}$ it is useful to consider the limit $\frac{z_2-z_1}{1+z_2} \ll 1$ and to assume that $\Phi(z_i)$ is fixed so its variation vanishes. Additionally, $\Phi_T \sim 1/5$ in our normalization. In this case we obtain $\frac{\delta\Phi_T(z)}{\Phi_T(z)} \simeq \epsilon \frac{z_2-z}{(1+z_2)}$. The relative change increases linearly, the maximal relative change being at z_1 , $\frac{\delta\Phi_T(z_1)}{\Phi_T(z_1)} \simeq \epsilon \frac{z_2-z_1}{(1+z_2)}$. The relative change is linear in the absolute change ϵ in w_{tot} and linear in the width $z_1 - z_2$ where the change occurs. Earlier changes in w_{tot} produce smaller changes in Φ_T . We now turn to the range $z < z_1$. Recall that for $z < z_1$ w_{tot} is unchanged, so it is better to choose $z_i = z_1$ in eq.(13). In this case $\delta I = 0$ and the only source of change is now the initial variation of $\Phi_T(z_1)$. Additionally, $I(z_1) = 0$. It follows from eq.(13) that $\delta\Phi_T(z) = e^{-I(z)} \delta\Phi_T(z_1)$. To get an idea about the numerical factors consider the case that w_{tot} is approximately constant, then $e^{-I(z)} \simeq \left(\frac{1+z}{1+z_1} \right)^{\frac{5+3w_{\text{tot}}}{2}}$, and $\frac{\delta\Phi_T(z)}{\Phi_T(z)} \simeq \epsilon \frac{z_2-z_1}{(1+z_2)} \left(\frac{1+z}{1+z_1} \right)^{\frac{5+3w_{\text{tot}}}{2}}$. Since $5/2 \gtrsim \frac{5+3w_{\text{tot}}}{2} \gtrsim 3/2$ for $z < z_1$, $\delta\Phi_T/\Phi_T$ slowly decreases from its maximal value at z_1 . Our estimates are in good agreement with the numerical results.

We can now estimate δJ , $\delta J(z) = \int_0^z dy \frac{1}{H(y)} \left(\frac{d_L(z)}{1+z} - \frac{d_L(y)}{1+y} \right) \Phi_T^2(y) \left[-\frac{\delta H(y)}{H(y)} + \frac{\delta d_L(z)}{d_L(z)} - \frac{\delta d_L(y)}{d_L(y)} + 2\frac{\delta\Phi_T(y)}{\Phi_T(y)} \right]$. Again, this fits well with the numerical estimates. We should emphasize an important feature. If δw_{tot} vanishes for values less than some redshift z_* then δd_L and δH vanish because they depend on an integral from $z = 0$ to z_* . In this case only $\delta\Phi_T$ is non-vanishing. For $z > z_*$ the situation reverses. Now $\delta\Phi_T$ vanishes, and the only contribution comes from the other terms. In Fig. 1 we show the results of the numerical evaluation of the induced relative differences in J due to modifications of w_{tot} of the form discussed above.

If the content of the universe is parametrized in terms of a cold matter component with relative density today of Ω_m and a DE component with a constant EOS parameter w_Q




FIG. 1: The relative changes in $J(z)$ induced by modifications in w_{tot} . In the top panel we show the EOS of the $\Omega_m = 0.3$, Λ CDM fiducial model and the EOS of three modified models. In the middle panel the relative deviations of the three modified EOS from the EOS of the fiducial model are shown, and in the lower panel the corresponding relative deviations of J .

and relative density $\Omega_Q = 1 - \Omega_m$, then the constraints on the function $w_{\text{tot}}(z)$ become constraints on the parameters Ω_m and w_Q . We discuss this example mainly to compare with the sensitivity of d_L measurements as is often done in the literature.

In Figure 2 we show 1% and 2% distinguishability contours of J (filled) and d_L (empty) in the range $z < 1.5$. The fiducial model is a $\Omega_m = 0.3$ Λ CDM model. Two models are deemed distinguishable at 1% if their J 's (d_L 's) deviate by more than 1% in the whole of the redshift range. Also shown are relative deviations of $J(z)$ from the fiducial model for four different constant EOS models. The d_L contours in Fig. 2 represent much better accuracy than current measurements [1, 2]. They are approximately equal to the final precision goals of the Supernovae Legacy Survey $1\sigma \sim 1\%$, $2\sigma \sim 2\%$ [35]. As can be seen from the figure a measurement of J with the same precision of a measurement of d_L gives about an order of magnitude improvement in determining the cosmological parameters. As emphasized previously, the figure should *not* be interpreted as a measurement of w_Q or of Ω_m , rather a measurement of the total EOS w_{tot} . Despite the fact that the observable J is related to




FIG. 2: The upper figure shows 1% and 2% distinguishability contours of J (filled) and d_L (empty) in the range $z < 1.5$. The fiducial model is a $\Omega_m = 0.3$ Λ CDM model. In the lower figure we show relative deviations of $J(z)$ from the fiducial model for four different constant EOS models labelled by (Ω_m, w_Q) : (from top to bottom) $(.32, -1.05)$, $(.31, -1.025)$, $(.29, -.975)$, $(.28, -.95)$.

perturbations in the matter only, and despite the fact that the J contours in Fig. 2 appear to be d_L contours with a limited range of Ω_m , Ω_m is not really measured separately here.

We have used a simplified model to obtain our sensitivity estimates to which many improvements and refinements can be made. We have found that obtaining a limiting accuracy as small as 10% over a range of redshift as small as 0.1 for local variations in w_{tot} seems plausible. The implementation of our method will require extraordinary observational efforts because it requires very accurate measurements of the shear, very accurate determination of redshifts of the sources, and in addition very accurate measurements of d_L as a function of redshift. To achieve such ambitious observational goals would require combining the results of several accurate experiments.

We believe that determining the nature of dark energy is still an outstanding problem. An independent determination of Ω_m (if it is possible to do so) will turn the constraints on w_{tot} into a constraints on the EOS of the DE. However, there remain degeneracies for groups of expansion histories that oscillate about each other [8]. This type of degeneracy

can perhaps be overcome by combining d_L and other observables with measurements of J . In any case, we do expect some remaining degeneracy that will have to be resolved by better theoretical understanding of the nature of dark energy.

This research was supported in part by the National Science Foundation under Grant No. PHY99-07949.

-
- [1] P. Astier *et al.*, “The Supernova Legacy Survey: Measurement of Ω_M , Ω_Λ and w from the First Year Data Set,” arXiv:astro-ph/0510447.
 - [2] A. G. Riess *et al.* [Supernova Search Team Collaboration], “Type Ia Supernova Discoveries at $z > 1$ From the Hubble Space Telescope: Evidence for Past Deceleration and Constraints on Dark Energy Evolution,” *Astrophys. J.* **607**, 665 (2004) [arXiv:astro-ph/0402512].
 - [3] D. N. Spergel *et al.*, “Wilkinson Microwave Anisotropy Probe (WMAP) three year results: Implications for cosmology,” arXiv:astro-ph/0603449.
 - [4] M. Tegmark *et al.*, “Cosmological Constraints from the SDSS Luminous Red Galaxies,” arXiv:astro-ph/0608632.
 - [5] W. J. Percival *et al.*, “Measuring the matter density using baryon oscillations in the SDSS,” arXiv:astro-ph/0608635.
 - [6] I. Maor, R. Brustein and P. J. Steinhardt, “Limitations in using luminosity distance to determine the equation-of-state of the universe,” *Phys. Rev. Lett.* **86**, 6 (2001) [Erratum-ibid. **87**, 049901 (2001)] [arXiv:astro-ph/0007297].
 - [7] J. Weller and A. Albrecht, “Opportunities for future supernova studies of cosmic acceleration,” *Phys. Rev. Lett.* **86**, 1939 (2001) [arXiv:astro-ph/0008314].
 - [8] I. Maor, R. Brustein, J. McMahon and P. J. Steinhardt, “Measuring the Equation-of-state of the Universe: Pitfalls and Prospects,” *Phys. Rev. D* **65**, 123003 (2002) [arXiv:astro-ph/0112526].
 - [9] J. A. Frieman, D. Huterer, E. V. Linder and M. S. Turner, “Probing dark energy with supernovae: Exploiting complementarity with the cosmic microwave background,” *Phys. Rev. D* **67**, 083505 (2003) [arXiv:astro-ph/0208100].
 - [10] P. S. Corasaniti and E. J. Copeland, “A model independent approach to the dark energy equation of state,” *Phys. Rev. D* **67**, 063521 (2003) [arXiv:astro-ph/0205544].

- [11] U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, “Exploring the Expanding Universe and Dark Energy using the Statefinder Diagnostic,” *Mon. Not. Roy. Astron. Soc.* **344**, 1057 (2003) [arXiv:astro-ph/0303009].
- [12] E. J. Copeland, M. Sami and S. Tsujikawa, “Dynamics of dark energy,” arXiv:hep-th/0603057.
- [13] A. Cooray, D. Huterer and D. Baumann, “Growth Rate of Large Scale Structure as a Powerful Probe of Dark Energy,” *Phys. Rev. D* **69**, 027301 (2004) [arXiv:astro-ph/0304268].
- [14] E. V. Linder, “Cosmic growth history and expansion history,” *Phys. Rev. D* **72**, 043529 (2005) [arXiv:astro-ph/0507263].
- [15] E. Bertschinger, “On the Growth of Perturbations as a Test of Dark Energy,” arXiv:astro-ph/0604485.
- [16] A. N. Taylor, T. D. Kitching, D. J. Bacon and A. F. Heavens, “Probing dark energy with the shear-ratio geometric test,” arXiv:astro-ph/0606416.
- [17] C. Heymans *et al.*, “Cosmological weak lensing with the HST GEMS survey,” *Mon. Not. Roy. Astron. Soc.* **361**, 160 (2005) [arXiv:astro-ph/0411324].
- [18] H. Hoekstra *et al.*, “First cosmic shear results from the Canada-France-Hawaii Telescope Wide Synoptic Legacy Survey,” arXiv:astro-ph/0511089.
- [19] E. Semboloni *et al.*, “Cosmic Shear Analysis with CFHTLS Deep data,” arXiv:astro-ph/0511090.
- [20] J. Albert *et al.* [SNAP Collaboration], “Probing Dark Energy via Weak Gravitational Lensing with the SuperNova Acceleration Probe (SNAP),” arXiv:astro-ph/0507460.
- [21] D. Huterer, “Weak Lensing and Dark Energy,” *Phys. Rev. D* **65**, 063001 (2002) [arXiv:astro-ph/0106399].
- [22] W. Hu, “Dark Energy and Matter Evolution from Lensing Tomography,” *Phys. Rev. D* **66**, 083515 (2002) [arXiv:astro-ph/0208093].
- [23] B. Jain and A. Taylor, “Cross-correlation Tomography: Measuring Dark Energy Evolution with Weak Lensing,” *Phys. Rev. Lett.* **91**, 141302 (2003) [arXiv:astro-ph/0306046].
- [24] F. Simpson and S. Bridle, “Illuminating Dark Energy with Cosmic Shear,” *Phys. Rev. D* **71**, 083501 (2005) [arXiv:astro-ph/0411673].
- [25] A. Upadhye, M. Ishak and P. J. Steinhardt, “Dynamical dark energy: Current constraints and forecasts,” *Phys. Rev. D* **72**, 063501 (2005) [arXiv:astro-ph/0411803].
- [26] C. Schimd *et al.*, “Tracking quintessence by cosmic shear: Constraints from VIRMOS-Descart

- and CFHTLS and future prospects,” arXiv:astro-ph/0603158.
- [27] L. Van Waerbeke and Y. Mellier, “Gravitational Lensing by Large Scale Structures: A Review,” arXiv:astro-ph/0305089.
 - [28] P. Schneider, “Weak Gravitational Lensing,” arXiv:astro-ph/0509252.
 - [29] D. Huterer and M. J. White, “Nulling Tomography with Weak Gravitational Lensing,” Phys. Rev. D **72**, 043002 (2005) [arXiv:astro-ph/0501451].
 - [30] D. Levy and R. Brustein, “Expressing the equation of state parameter in terms of the three dimensional cosmic shear,” arXiv:astro-ph/0607435.
 - [31] V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, “Theory Of Cosmological Perturbations,” Phys. Rept. **215**, 203 (1992).
 - [32] C. P. Ma and E. Bertschinger, “Cosmological perturbation theory in the synchronous and conformal Newtonian gauges,” Astrophys. J. **455**, 7 (1995) [arXiv:astro-ph/9506072].
 - [33] D. Levy and R. Brustein, to appear.
 - [34] M. Zaldarriaga and U. Seljak, “An All-Sky Analysis of Polarization in the Microwave Background,” Phys. Rev. D **55**, 1830 (1997) [arXiv:astro-ph/9609170].
 - [35] <http://www.cfht.hawaii.edu/SNLS/>